First Principles Study on the Stability and Electronic Structures of 7,8-Dichloro-4-Oxo-4H-Chromene-3-Carbaldehyde

Pek-Lan Toh^{1,a,*}, Montha Meepriruk^{2,b}, Lee Sin Ang^{3,c}, Shukri Sulaiman^{4,d} and Mohamed Ismail Mohamed-Ibrahim^{4,e}

¹Department of Electronic Engineering, Faculty of Engineering and Green Technology, Universiti Tunku Abdul Rahman, 31900 Perak, Malaysia

²School of Chemistry, Institute of Science and Technology, Kamphaeng Phet Rajabhat University, Kamphaeng Phet 62000, Thailand

³Faculty of Applied Sciences, Universiti Teknologi Mara, 02600 Perlis, Malaysia

⁴Computational Chemistry and Physics Laboratory, School of Distance Education, Universiti Sains Malaysia, 11800 Penang, Malaysia

^apeklan_toh@yahoo.com.my, ^bmontha_mee@hotmail.com, ^canglee631@perlis.uitm.edu.my, ^dshukri@usm.my, ^emi-mi@usm.my

Keywords: Density Functional Theory (DFT), 7,8-Dichloro-4-Oxo-4H-Chromene-3-Carbaldehyde, geometric parameters, electronic structures, Mulliken atomic charges.

Abstract. The molecular structures and electronic properties of 7,8-Dichloro-4-Oxo-4H-Chromene-3-Carbaldehyde, $C_{10}H_4Cl_2O_3$ have been studied using Density Functional Theory (DFT) method. The calculation of geometry optimization was conducted to find the local energy minimum of $C_{10}H_4Cl_2O_3$ molecular system. The equilibrium geometries were used to determine the HOMO-LUMO gaps, Mulliken atomic charges, and other electronic structures of $C_{10}H_4Cl_2O_3$. The significant findings from DFT/B3LYP functional within the basis sets of 6-31G**, 6-31++G**, 6-311G**, and 6-311++G** show that the optimized geometries of $C_{10}H_4Cl_2O_3$ are in good agreement with that of measurement data. To further investigate this, using a variety of basis sets (3-21G, 6-31G, 6-31++G, 6-31G**, 6-31++G**, 6-311G, 6-311++G, 6-311G**, and 6-311++G**), it is found that the calculated total energy values of $C_{10}H_4Cl_2O_3$ are close to each other. Similarly, the computed HOMO-LUMO energy gaps obtained are also close to each other. Using the scheme of Mulliken Population Analysis (MPA), the trend of findings are the same for both cases of B3LYP/6-31G and B3LYP/6-31G** level of calculations. For the method of B3LYP/6-31G, it is clearly found that C_4 and C_6 have the highest positively charge, with the corresponding values about +0.284 and +0.238, respectively. On the other hand, the charge values of +0.157 and +0.206 are found on Cl_1 and Cl_2 atoms. The atoms of O_1 , O_2 , and O_3 have the negatively charges, with the values of about 0.398, -0.512, and -0.424, respectively. Similarly, in the case of DFT/B3LYP/6-31G** level of theory, the computed charge values of C_2 and C_4 are about +0.311 and +0.393, respectively. Furthermore, the Cl_1 and Cl_2 atoms have the positively charge values of about +0.043 and +0.070, whereas for the O_1 , O_2 , and O_3 atoms, the charges values obtained are about -0.421, -0.467, and -0.498, respectively.

Introduction

Many substituted 3-Formylchromone has been synthesized, and their properties have been studied [1-6]. There have been recently received considerable attentions as a subject to the academic researches. Several 3-Formylchromone derivatives could also serve as key synthetic building blocks for the molecular construction in the biological fields and various medical chemistry applications primarily [1-6]. In the literature, a number of experimental studies concerning 3-Formylchromone derivatives have been reported. In 2014, Parveen *et al.* characterized a series of new substituted 3-formylchromone derivatives using Knoevenagel Condensation (KC) technique [1]. They also worked on Mass Spectrometry (MS), Infrared (IR), and Raman spectroscopy studies of these new substituted 3-formylchromone. The experimental data clearly indicates that these synthesized

3-formylchromone derivatives bind strongly with DNA molecules. The structures of substituted 3-formylchromone obtained are quite similar to bis-intercalators of DNA, which can be reserved as a template for future studies in the therapeutic purposes. In 2015, Ishikawa also studied the new crystal structure and characterization of 7,8-Dichloro-4-Oxo-4H-Chromene-3-Carbaldehyde, $C_{10}H_4Cl_2O_3$ using X-ray crystallography, IR, Raman spectroscopy, and others [6]. Density Functional Theory (DFT) calculations nowadays have been widely used as powerful method to investigate geometries and electronic properties of various materials [7-13]. Babar *et al.*, in 2014, investigated the Highest Occupied Molecular Orbital (HOMO) and Lowest Unoccupied Molecular Orbital (LUMO) energies of a series of new substituted 2-anilinothiazoles derivatives using the DFT/B3LYP/6-31G* level of theory [7]. The calculation results found that the 2-anilinothiazoles substitutions have no significant effect on the HOMO-LUMO energy gaps. In 2015, Singh reported the effect on geometrical parameters, atomic charges, and vibrational frequencies of uracil and 5-Methyuracil (Thymine) molecules [9]. The computed vibrational frequencies obtained are in good agreement with experimental measurements. Also, Toh *et al.*, in 2015, used the DFT method employing the three functionals with the 6-31++G** and 6-311++G** basis sets to investigate the electronic structures, and hyperfine interactions of muonium in imidazole and 1-Methylimidazole molecular system, respectively [11]. The results found that the calculated total energy values obtained of both muoniated systems are consistent with measurement data. Furthermore, there is no reported computational investigation of 7,8-Dichloro-4-Oxo-4H-Chromene-3-Carbaldehyde, $C_{10}H_4Cl_2O_3$ compound in the literature. To the best of our knowledge, first principles DFT procedure was carried out to study the geometric parameters, local energy minima, Mulliken atomic charges, and others for the $C_{10}H_4Cl_2O_3$ molecular system. The overall details of the computational methodology, results, discussion, and summary are provided in the following sections.

Computational Methodology

All Density Functional Theory (DFT) calculations, including geometry optimization and single point calculation in this work were carried out using the Density Functional Theory (DFT) procedure within Gaussian 03 and Gaussian 09 program packages [14,15]. A single molecule of 7,8-Dichloro-4-Oxo-4H-Chromene-3-Carbaldehyde, $C_{10}H_4Cl_2O_3$ was chosen to simulate the local host environment. The sketch of molecular structure of the $C_{10}H_4Cl_2O_3$ with atom numbering is illustrated in Fig. 1. Using the $C_{10}H_4Cl_2O_3$ molecular system, geometry optimization calculations at the DFT/B3LYP level of theory with 3-21G, 6-31G, 6-31++G, 6-31G**, 6-31++G**, 6-311G, 6-311++G, 6-311G**, and 6-311++G** basis sets were performed to find the local energy minimum of $C_{10}H_4Cl_2O_3$. The optimized geometries of $C_{10}H_4Cl_2O_3$ were then used to determine the total energy and Frontier molecular orbital energies, particularly the Highest Occupied Molecular Orbital (HOMO) and Lowest Unoccupied Molecular Orbital (LUMO) of $C_{10}H_4Cl_2O_3$. Using the Mulliken population analysis (MPA), the atomic charges of $C_{10}H_4Cl_2O_3$ were also calculated.

Fig. 1: The numbering system used for 7,8-Dichloro-4-Oxo-4H-Chromene-3-Carbaldehyde, $C_{10}H_4Cl_2O_3$.

Results and Discussion

The optimized molecular geometries of 7,8-Dichloro-4-Oxo-4H-Chromene-3-Carbaldehyde, $C_{10}H_4Cl_2O_3$ are investigated by DFT/B3LYP method using several basis sets of 3-21G, 6-31G, 6-31++G, 6-31G**, 6-31++G**, 6-311G, 6-311++G, 6-311G**, and 6-311++G**. In this work, the calculated optimized bond lengths, bond angles, and dihedral angles obtained for the title compound are compared with the experimental values and summarized in Table 1. It can be clearly observed from the table that the optimized bond lengths at 6-31G**, 6-31++G**, 6-311G**, and 6-311++G** basis sets are in rather good agreement with the experimental data, which reported by Ishikawa [6]. The overall computational data are only different with those of experimental values in the range from 0.01% to 2.5%. While for the other basis sets of 3-21G, 6-31G, 6-31++G, 6-311G, and 6-311++G, the calculated bond lengths obtained are slightly different from the experimental ones for a range of 0.1% - 5.0%. Furthermore, the double bond of C_1-O_1 is the shortest bond length among the bonds in $C_{10}H_4Cl_2O_3$. While the bond lengths of C_9-Cl_1 and C_9-Cl_2 have the longest values in the title molecule, and the order of these C-Cl bonds are equal to 1. In the case of bond angles, the computed values obtained are close to that of experimental results, which given by Ishikawa [6]. Using DFT/B3LYP simulation approach within a variety of basis sets (3-21G, 6-31G, 6-31++G, 6-31G**, 6-31++G**, 6-311G, 6-311++G, 6-311G**, and 6-311++G**), the calculated bond angles are only in the range of 0.1% - 1.1% different with those of measurement values [6]. For the dihedral angles of $C_{10}H_4Cl_2O_3$, the computed values at the DFT/B3LYP level within a variety of basis sets are found to vary in the range of $2.93^\circ - 3.14^\circ$, which are slightly smaller to the measurement data [6].

Table 1: Selected optimized geometrical parameters of $C_{10}H_4Cl_2O_3$ molecular system.

	3-21G	6-31G	6-31++G	6-31G**	6-31G**	6-311G	6-311++G	6-311G**	6-311++G**	Experiment [6]
Bond Length (Angstrom)										
C_1-O_1	1.237	1.244	1.246	1.218	1.220	1.241	1.242	1.211	1.212	1.207
C_1-C_2	1.477	1.477	1.478	1.483	1.484	1.476	1.476	1.485	1.485	1.484
C_3-O_2	1.373	1.367	1.366	1.342	1.342	1.367	1.365	1.340	1.339	1.346
C_4-O_2	1.393	1.396	1.395	1.372	1.373	1.396	1.395	1.370	1.370	1.374
C_6-O_3	1.244	1.254	1.255	1.228	1.230	1.253	1.253	1.221	1.223	1.233
C_9-C_{10}	1.383	1.393	1.393	1.398	1.399	1.389	1.390	1.394	1.394	1.387
C_9-Cl_1	1.809	1.805	1.804	1.741	1.741	1.806	1.806	1.741	1.740	1.723
$C_{10}-Cl_2$	1.798	1.795	1.795	1.735	1.735	1.797	1.797	1.735	1.734	1.723
Bond Angle (Degree)										
$C_2-C_1-O_1$	123.35	122.96	122.93	123.50	123.43	122.90	122.85	123.65	123.56	123.78
$C_3-O_2-C_4$	118.23	119.21	119.41	119.04	119.26	119.29	119.43	119.03	119.17	118.30
$C_2-C_6-O_3$	123.63	122.95	122.82	123.53	123.38	122.87	122.83	123.58	123.50	123.48
$C_5-C_6-O_3$	122.80	122.11	122.01	122.72	122.60	122.00	121.96	122.77	122.68	122.34
$C_3-C_9-Cl_1$	118.30	118.11	118.14	118.51	118.54	118.08	118.12	118.51	118.53	119.33
$C_{10}-C_9-Cl_1$	120.41	120.78	120.78	120.68	120.69	120.74	120.74	120.68	120.69	119.94
$C_9-C_{10}-Cl_2$	122.46	122.45	122.37	122.37	122.32	122.46	122.41	122.34	122.34	122.86
$C_4-C_{10}-Cl_2$	118.49	119.30	119.38	119.50	119.58	119.32	119.34	119.46	119.50	118.83
Dihedral Angle (Degree)										
$Cl_2-C_{10}-C_4-O_2$	2.93	3.09	3.13	3.07	3.08	3.14	3.19	3.09	3.11	3.15

In Table 2, the total energies and HOMO-LUMO gaps of $C_{10}H_4Cl_2O_3$ at the DFT/B3LYP level within a variety of basis sets (3-21G, 6-31G, 6-31++G, 6-31G**, 6-31++G**, 6-311G, 6-311++G, 6-311G**, and 6-311++G**) are presented. According to the data from the table, the computed total energies of $C_{10}H_4Cl_2O_3$ molecular system are calculated to be in the range from -41408.704 eV

to -41626.097 eV. As the value of -41626.097 eV obtained from the basis set of 6-311++G** are chosen to be a zero energy, the total energy values of the other basis sets (3-21G, 6-31G, 6-31++G, 6-31G**, 6-31++G**, 6-311G, 6-311++G, and 6-311G**) are only 0.01% - 1.18% greater than the DFT/B3LYP/6-311++G** simulation approach. In addition, the HOMO-LUMO energy gaps of $C_{10}H_4Cl_2O_3$ molecular system are also presented in Table 2. Using DFT/B3LYP computational calculations in a variety of basis sets (3-21G, 6-31G, 6-31++G, 6-31G**, 6-31++G**, 6-311G, 6-311++G, 6-311G**, and 6-311++G**), the HOMO-LUMO energy gaps fall in the range of 4.21 eV - 4.64 eV. Also, it can be noted that the energy gaps of all the basis set are close to each other in this study. As shown in Fig. 2, the molecular orbital diagrams of the frontier orbitals obtained in the study for the title molecule are illustrated using the DFT/B3LYP functional calculation employing the basis set of 6-311G**. For $C_{10}H_4Cl_2O_3$ molecular system, the HOMO has mostly p character, respectively, comes from the p_x orbitals of the O_1 , O_3 , C_1 , C_2 , and C_6 . In other words, the electron distributions of the HOMO diagram are mainly localized on the parts of aldehyde and C_4-O_2 , respectively. The results, given in the figure, also presented that the LUMO has mostly p character symmetry, where the electron distributions are localized on the whole molecular system.

Table 2: Calculated total and frontier molecular orbital energies (eV) of $C_{10}H_4Cl_2O_3$.

	3-21G	6-31G	6-31++G	6-31G**	6-31G**	6-311G	6-311++G	6-311G**	6-311++G**
Total energy	-41133.29	-41408.70	-41614.11	-41614.78	-41620.30	-41620.94	-41619.72	-41620.14	-41625.73
HOMO	-4.24	-7.03	-7.24	-7.56	-7.01	-7.37	-7.48	-7.59	-7.25
LUMO	-0.02	-2.51	-2.75	-3.07	-2.38	-2.76	-2.97	-3.09	-2.61
HOMO-LUMO gap	4.21	4.52	4.49	4.49	4.62	4.61	4.51	4.50	4.64

Fig. 2: Calculated molecular orbital diagrams of the frontier orbitals of $C_{10}H_4Cl_2O_3$ at the DFT/B3LYP/6-311G** level of theory.

Using Mulliken Population Analysis (MPA) at B3LYP/6-31G and B3LYP/6-31G** level of calculations, the atomic charge distributions of $C_{10}H_4Cl_2O_3$ are illustrated in Fig. 3. The results in the case of DFT/B3LYP functional calculations employing with 6-31G basis set, given in Fig. 3(a), clearly indicate that C_4 atom has the highest positively charge, with the corresponding value of charge about +0.284. By following the charge of C_6 is determined to be +0.238 in $C_{10}H_4Cl_2O_3$ molecular system. Furthermore, the computed positively charge of Cl_1 and Cl_2 atoms, respectively are about +0.157 and +0.206. While for the oxygen, O_1 , O_2 , and O_3 atoms become negatively charges, the charges are calculated to be about -0.398, -0.512, and -0.424, respectively. In this study, the computational data evident that Mulliken atomic charge distributions are dependent on the size of basis sets. However, for both B3LYP/6-31G and B3LYP/ 6-31G** calculations, the trend of results are the same. In the case of DFT/ B3LYP/ 6-31G** level of theory (Fig. 3(b)), both atoms of C_4 and C_6 becomes more positive than B3LYP/6-31G calculation, respectively, with the corresponding values about +0.311 and +0.393. For the title molecule, the charges of Cl_1 and Cl_2 are

then determined to be about +0.043 and +0.070, whereas the charges values of -0.421, -0.467, and -0.498 are obtained on the O₁, O₂, and O₃ atoms, respectively.

Fig. 3: Mulliken atomic charges distribution used for the optimized C₁₀H₄Cl₂O₃ based on the (a) B3LYP/6-31G and (b) B3LYP/6-31G** level of calculations.

Summary

All computational calculations were carried out using the Gaussian 03 and Gaussian 09 program packages [14,15] within first principles investigation within the Density Functional Theory (DFT) method. In the case of geometrical parameters of C₁₀H₄Cl₂O₃ molecular system, the computed values are close to experimental measurements, which presented by Ishikawa [6]. In addition, the calculated total energies are close to each other using DFT/B3LYP level of calculation within a variety of basis sets (3-21G, 6-31G, 6-31++G, 6-31G**, 6-31++G**, 6-311G, 6-311++G, 6-311G**, and 6-311++G**). Similarly, the HOMO-LUMO energy gaps of title compound obtained are consistent with each other. In the case of Mulliken atomic charges, the computed charges of C₄ and C₆ atoms are about +0.284 and +0.238, which have the highest positively charges in the title compound using DFT/B3LYP/6-31G simulation approach. The charges of Cl₁ and Cl₂ atoms are found to be about +0.157 and +0.206, while the charge values of -0.398, -0.512, and -0.424, respectively, are obtained on the atoms of O₁, O₂, and O₃. The similar trend is found in the case of DFT/B3LYP/6-31G** calculation. The charge values of C₄ and C₆ atoms are about +0.311 and +0.393. By following the charges of Cl₁ and Cl₂ atoms are calculated to be about +0.043 and +0.070, whereas the negatively charge values of -0.421, -0.467, and -0.498 are found for the O₁, O₂, and O₃ atoms, respectively. Further studies are currently being carried out to investigate the effects of cluster size in C₁₀H₄Cl₂O₃ molecular system.

References

- [1] M. Parveen, A.M. Malla, Z. Yaseen, A. Ali, M. Alam, Synthesis, characterization, DNA-binding studies and Acetylcholinesterase inhibition activity of new 3-Formyl chromone derivatives, *J. Photochemistry and Photobiology B: Bio.* 130 (2014) 179-187.
- [2] Y. Ishikawa, Crystal structure of 6-Fluoro-4-Oxo-4H-Chromene-3-Carbaldehyde, *Acta Cryst. E.* 70 (2014) o774.
- [3] Y. Ishikawa, Crystal structure of 7-Chloro-4-Oxo-4H-Chromene-3-Carbaldehyde, *Acta Cryst. E.* 70 (2014) o831.
- [4] Y. Ishikawa, Crystal structure of 7-Bromo-4-Oxo-4H-Chromene-3-Carbaldehyde, *Acta Cryst. E.* 70 (2014) o996.

-
- [5] Y. Ishikawa, Crystal structure of 6-Bromo-7-Fluoro-4-Oxo-4H-Chromene-3-Carbaldehyde, *Acta Cryst. E.* 71 (2015) 857-860.
- [6] Y. Ishikawa, Crystal structure of 7,8-Dichloro-4-Oxo-4H-Chromene-3-Carbaldehyde, *Acta Cryst. E.* 71 (2015) 902-905.
- [7] A. Babar, H. Khalid, K. Ayub, S. Saleem, A. Waseem, T. Mahmood, M.A. Munawar, G. Abbas, A.F.Khan, Synthesis, characterization and Density Functional Theory study of some new 2-Anilinothiazoles, *J. Molecular Structure* 1072 (2014) 221-227.
- [8] J.S. Singh, FT-IR and Raman spectra, ab initio and density functional computations of the vibrational spectra, molecular geometries and atomic charges of uracil and 5-Halogenated uracils (5-X-Uracils; X = F, Cl, Br, I), *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy* 117 (2013) 502-518.
- [9] J.S. Singh, FT-IR and Raman spectra, ab initio and density functional computations of the vibrational spectra, molecular geometries and atomic charges of uracil and 5-Methyluracil (Thymine), *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy* 137 (2015) 625-640.
- [10] P.L. Toh, S. Sulaiman, M.I. Mohamed-Ibrahim, Computational studies of electronic structures and hyperfine interactions of muonium in tetraphenylgermane, *Advanced Materials Research* 630 (2013) 418-423.
- [11] P.L. Toh, S. Sulaiman, M.I. Mohamed-Ibrahim, L.S. Ang, Density functional theory studies of electronic structures and hyperfine interactions of muonium in imidazole, *Applied Mechanics and Materials* 749 (2015) 134-138.
- [12] P.L. Toh, M. Meepripruk, L.S. Ang, S. Sulaiman, M.I. Mohamed-Ibrahim, Density Functional Theory investigations on the geometric and electronic structures of 4-Azidomethyl-6-Isopropyl-2H-Chromen-2-One, *MATEC Web of Conferences* 27 (2015) 01003:1-5.
- [13] P.L. Toh, R. Rasmi, M. Meepripruk, L.S. Ang, S. Sulaiman, M.I. Mohamed-Ibrahim, First principles Density Functional Theory investigation on the structural, energetic, and electronic properties of 6-Bromo-4-Oxo-4H-Chromene-3-Carbaldehyde, *Applied Mechanics and Materials* 835 (2016) 308-314.
- [14] M.J. Frisch, G.W. Trucks, H.B. Schlegel, G.E. Scuseria, M.A. Robb, J.R. Cheeseman, *et al.*, Gaussian 03, Revision C.02, Gaussian, Inc. 2004.
- [15] M.J. Frisch, G.W. Trucks, H.B. Schlegel, G.E. Scuseria, M.A. Robb, J.R. Cheeseman, *et al.*, Gaussian 09, Revision D.01, Gaussian, Inc. 2009