Influence of Organic Supplements on Growth and Development of *In Vitro* Shoots of *Bulbophyllum dhaninivatii* Seidenf

Anupan Kongbangkerd^{1,a*}, Santi Watthana^{2,b} and Kanok-Orn Srimuang^{3,c}

¹Plant Tissue Culture Research Unit, Department of Biology, Faculty of Science, Naresuan University, Phitsanulok 65000 Thailand

²School of Biology, Institute of Science, Suranaree University of Technology, Nakhon Ratchasima 30000 Thailand

³Department of Biotechnology, Faculty of Agriculture & Natural Resources, University of Payao, Payao 56000 Thailand

^{a*}anupank@nu.ac.th, ^bsantiqsb@gmail.com, ^cs.kanokorn@gmail.com

Keywords. *Bulbophyllum*, Growth, Organic supplements

Abstract. *In vitro* shoot culture of *Bulbophyllum dhaninivatii* Seidenf. was conducted on semi-solid Vacin and Went (1949) medium supplemented with coconut water (50 100 and 150 ml/L), potato extract (25 and 50 g/L) and banana homogenate (25 and 50 g/L) for 12 weeks. The results found that the highest shoot number (6.92 shoots) could observe on the medium with 150 ml/L coconut water, 50 g/L potato extract and 50 g/L banana homogenate while the highest leaf and root number could obtain when cultured on the medium supplemented with 100 ml/L coconut water, 50 g/L potato extract and 50 g/L banana homogenate.

Introduction

Bulbophyllum is one of the largest genera in the Orchidaceae family. This genus comprised of about 3,000 species [1]. In Thailand around 210 species of the *Bulbophyllum* genus are now identified [2]. *Bulbophyllum dhaninivatii* Seidenf. was firstly found at Phu Miang Sub-District, Chatrakarn District, Phitsanulok Province. Nowadays, this species is also distributed in Phu Luang Wild Life Sanctuary, Loei Province and listed as an endemic species of Thailand [3]. Because of a limited distribution area and specific habitat, this species seems to become risk of extinction from natural site. *In vitro* propagation techniques have been widely used for conservation and succeeded in many orchid species i.e. in *Dendrobium* [4-7] and *Cymbidium* species [8-10] etc. *In vitro* culture and micropropagation *Bulbophyllum* species have been reported [11-14] and factors improving shoot multiplication in *Bulbophyllum* species, especially medium components, were also studied [15-17]. However, influence of other medium components, particularly organic supplements, on shoot regeneration efficiency of *Bulbophyllum* species has only few performed specifically in *B. dhaninivatii*. The objective of this study was conducted to test the effects of different concentration and combination of coconut water, potato extract and banana homogenate on shoot proliferation and plantlet regeneration potentials of *B. dhaninivatii in vitro*. Our receiving results will be useful for conservation planning and used as a model for *in vitro* conservation of other *Bulbophyllum* species.

Materials and methods

In vitro shoots of *B. dhaninivatii* Seidenf. from stock culture were transferred to culture on semi-solid Vacin & Went medium [18] supplemented with 20 g/L sucrose, 2.0 g/l activated charcoal and 7.5 g/L agar. A different concentrations of a combination between coconut water (CW) (50, 100, 150 mL/L), potato extract (PE) (25, 50 g/L) and banana homogenate (BH) (25, 50 g/L) were additionally filled in the medium. The pH of medium was adjusted to 5.2. The cultures were placed under 12 hrs photoperiod of LED warm white light with 20 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$ PPF at 25-27 °C. The data were collected at 12 weeks after inoculation for number of shoots, roots and leaves, shoot, root and leave length, width of leaves. Experimental design was conducted on 3x2x2 Factorial in Complete

Randomize Design. Each treatment consisted of 15 replicates. The experiment was repeated thrice. Data collected were subjected to analysis of variance and means were separated by Duncan's Multiple Range Test (DMRT).

Results and discussions

An organic supplements, one of an important medium component, have been used to stimulate growth, development and promote shoot regeneration in many orchid tissue cultures i.e. in *Dendrobium* [19], *Cymbidium pendulum* [20], *Vanda* and *Mokara* [21], *Vanda helvola* [22] and *Cypripedium macranthos* [23] etc. *In vitro* shoots of *B.dhaninivatii* were cultured on semi-solid VW medium supplemented with different concentrations of a combination between CW, PE and BH for 12 weeks. The results indicated that morphological change of *in vitro* shoots could be noticed after the second week of culture. Enlargements of explants with shoot bud induction as well as new root formation in all treatments were then observed after 4 weeks of culture. New regenerated shoots with green leaves were then investigated at 8 weeks and continuously grew into plantlets after 12 weeks of culture. The overall results found that adding organic supplements to the medium improved better growth and morphological change of *in vitro* shoots than that cultured on the control medium. The highest shoot regeneration number was observed on the medium supplemented with highest concentration of CW (150 mL/L), PE (50 g/L) and BH (50 g/L) (Table 1, Fig.1). No surprising results was noticed on this condition because CW contains several types of cytokinin [24] which could be promoted shoot multiplication in many orchids [25-26] included in *B.dhaninivatii* while PE and BH could also stimulate shoot growth and regeneration efficiency when combined with CW [20, 27]. The highest root and leaf number were obtained when maintaining the higher concentration of PE (50 g/L) and BH (50 g/L) with a reduction of CW to 100 mL/L. The related result has been observed in *Calanthe* hybrids [28]. In consideration of shoot, root and leaf growth in *B.dhaninivatii*, adding 50 mL/L CW with 25 g/L PE and 50 g/L BH to the medium promoted the highest shoot (3.59 cm) and root length (1.83 cm) as well as leaf width (0.58 cm) and length (2.87 cm), respectively. The relative effect of the suitable ratio between CW: PE: BH has been studied in *Vanda teres* [29], *Spathoglottis kimballiana* [30] and *Cymbidium pendulum* [20] etc.

Fig.1 Plantlet after cultured for 12 weeks. (A) VW medium. (B) VW medium supplemented with 50 mL/L CW, 25 g/L PE and 50 g/L BH. (C) VW medium supplemented with 100 mL/L CW, 50 g/L PE and 50 g/L BH. (D) VW medium supplemented with 150 mL/L CW, 50 g/L PE and 50 g/L BH.

Table 1. Effect of coconut water (CW), potato extract (PE) and banana homogenate (BH) supplementation on growth of *in vitro* shoot of *Bulbophyllum dhaninivatii* after cultured for 12 weeks.

CW [mL/L]	PE [g/L]	BH [g/L]	Shoots		Leaves			Roots		
			Number	Length [cm]	Number	Length [cm]	Width [cm]	Number	Length [cm]	
0	0	0	0.8 ±	1.26 ±	1.0 ±	0.73 ±	0.10 ±	3.1 ±	0.55 ±	
			0.2 b*	0.30 d	0.3 d	0.23 d	0.03 c	0.8 c	0.13 d	
			4.5 ±	2.19 ±	2.4 ±	2.52 ±	0.53 ±	7.3 ±	1.31 ±	
50	25	25	1.4 ab	0.44bcd	0.3 ab	0.48 ab	0.25 ab	1.2 abc	0.20 abc	
			4.1 ±	3.59 ±	2.2 ±	2.87 ±	0.58 ±	10.7 ±	1.83 ±	
			50	1.5 ab	0.52 a	0.3 abc	0.37 a	0.19 a	2.1 ab	0.22 a
	50	25	4.1 ±	3.50 ±	2.7 ±	2.74 ±	0.27 ±	9.5 ±	0.96 ±	
			1.5 ab	0.39a	0.4 ab	0.29 a	0.02 bc	1.5 ab	0.17 bcd	
			3.8 ±	3.07 ±	2.2 ±	2.45 ±	0.29 ±	9.1 ±	1.34 ±	
	50	50	2.1 ab	0.36abc	0.3 abc	0.31 abc	0.02 abc	1.3 ab	0.18 abc	
			2.6 ±	3.15 ±	2.8 ±	2.39 ±	0.29 ±	8.4 ±	1.53 ±	
			100	0.5 b	0.44 ab	0.3 a	0.36 abc	0.03 abc	0.8 ab	0.17 ab
100	25	25	2.1	1.97 ±	2.0 ±	1.53 ±	0.24 ±	9.3 ±	1.45 ±	
			50	±0.4 b	0.38cd	0.3 abcd	0.33bcd	0.03 bc	1.2 ab	0.15 abc
			3.0 ±	2.04 ±	2.5 ±	1.36 ±	0.20 ±	8.3 ±	1.04 ±	
	50	25	1.1 b	0.41bcd	0.6 ab	0.34 d	0.04 c	1.4 ab	0.11 bcd	
			2.3 ±	3.55 ±	2.9 ±	2.80 ±	0.34 ±	12.5 ±	1.33 ±	
			50	0.6 b	0.38 a	0.3 a	0.26 a	0.03 abc	1.5 a	0.21 abc
	50	50	3.0 ±	1.68 ±	1.5 ±	1.19 ±	0.17 ±	6.6 ±	1.03 ±	
			150	0.8 b	0.36 d	0.4 bcd	0.32 d	0.04 c	1.4 bc	0.20 bcd
			3.0 ±	1.53 ±	1.7 ±	0.97 ±	0.15 ±	10.6 ±	0.93 ±	
150	25	25	50	0.4 b	0.22 d	0.4 abcd	0.29 d	0.04 c	2.3 ab	0.13 cd
			4.4 ±	1.95 ±	1.7 ±	1.42 ±	0.24 ±	10.1 ±	1.17 ±	
			50	1.0 ab	0.35 cd	0.4 abcd	0.34 cd	0.06 bc	2.8 ab	0.19 bc
	50	25	7.0 ±	1.21 ±	1.1 ±	0.98 ±	0.15 ±	9.5 ±	1.24 ±	
			50	1.4 a	0.16 d	0.3 cd	0.27 d	0.04 c	1.6 ab	0.22 bc

*Values are means ± SD of 3 repeats (10 *in vitro* shoots per repeat). Different letters within the same column show highly significant differences analyzed by Duncan's new multiple range tests at $p \leq 0.05$.

Summary

From our investigation, the effects of different organic supplements on shoot regeneration and multiplication of *B.dhaninivatii* were studied and improved. The results revealed that adding 100 ml/L coconut water with 50 g/L potato extract and 50 g/L banana homogenate to the VW medium was the most influential for shoot regeneration and plantlet development. Organic supplements could promote shoot multiplication of *B.dhaninivatii* when compared to the control. Our study is an alternative way for rapid propagation of *B.dhaninivatii* without using plant growth regulators.

Acknowledgement

The author thanks to Naresuan University for providing research fund of the fiscal year 2016. We would like to acknowledge Department of Biology, Faculty of Science, Naresuan University for providing the research facilities in this study.

References

- [1] E.S. Siegerist, *Bulbophyllums and their allies: a grower's guide*, Timber Press, Inc., Portland, Oregon, 2001.
- [2] Information on <http://apps.kew.org/wcsp/advsearch.do;jsessionid=123D10881BDE409A73896025EA7EC668>
- [3] O. Thaithong, *Orchids of Thailand*, Office of Environmental Policy and Planning, Bangkok, Thailand, 1999.
- [4] M.M. Hossain, M. Sharma, P. Pathak, *In vitro* propagation of *Dendrobium aphyllum* (Orchidaceae)-seed germination to flowering. *J. Plant Biochem. Biotechnol.* 22(2) (2013) 157–167.
- [5] D. Zhao, G. Hu, Z. Chen, Y. Shi, L. Zheng, A. Tang, C. Long, Micropropagation and *in vitro* flowering of *Dendrobium wangliangii*: a critically endangered medicinal orchid. *J. Med. Plants Res.* 7(28) (2013) 2098-2110.
- [6] S. Parthibhan, M.V. Rao, T.S. Kumar, *In vitro* regeneration from protocorms in *Dendrobium aqueum* Lindley – an imperiled orchid. *J. Gen. Eng. Biotechnol.* 13 (2015) 227-233.
- [7] J.A.T. da Silva, J.C. Cardoso, J. Dobra'nszki, S. Zeng, *Dendrobium* micropropagation: a review. *Plant Cell Rep.* 34 (2015) 671-704.
- [8] J.A.T. da Silva, M. Tanaka, Culture vessel affects hybrid *Cymbidium* protocorm-like body and callus induction. *Flor. Ornament. Biotechnol.* 3(1) (2009) 53-55.
- [9] J.A.T. da Silva, New basal media for protocorm-like body and callus induction of hybrid *Cymbidium*. *J. Fruit Ornament. Plant Res.* 20(2) (2012) 127-133.
- [10] P. Mohanty, S. Paul, M.C. Das, S. Kumaria, P. Tandon, A simple and efficient protocol for the mass propagation of *Cymbidium mastersii*: an ornamental orchid of Northeast India. *AoB PLANTS* 2012: pls023; doi:10.1093/aobpla/pls023
- [11] S.K. Bhadra, H. Barua, M.M. Hossain, *In vitro* germination and rapid micropropagation of *Bulbophyllum lilacinum* Redley. *Bangladesh J. Bot.* 33(2) (2004)103-107.
- [12] Y.I. Lee, E. Yeung, Embryo development and *in vitro* seed germination of *Bulbophyllum fascinator*. *Acta Hort.* 878 (2010) 243-250.
- [13] C. Maneerattananarungroj, S. Laywisadkul, A. Kongbangkerd, Tissue culture of *Bulbophyllum affine* Lindl. *NU Sci. J.* 7(2) (2011) 45-59.
- [14] M.M.M. Than, *In vitro* conservation of endangered orchid *Bulbophyllum auricomum* Lindl. *Prop. Ornament. Plants* 13(4) (2013) 154-159.
- [15] S.P. Vij, A. Kher, P. Pathak, Regeneration competence of *Bulbophyllum careyanum* (Hook.) Spreng. pseudobulb segments. *J. Orchid Soc. India* 14 (2000) 47–55.
- [16] M.M.M. Than, A. Pal, S. Jha, *In vitro* flowering and propagation of *Bulbophyllum auricomum* Lindl., the royal flower of Myanmar. *Acta Hort.* 829 (2009) 105-111.
- [17] M.M.M. Than, A. Pal, S. Jha, Plant regeneration from callus cultures in endangered orchid *Bulbophyllum auricomum* Lindl. *Prop. Ornament. Plants*, 12(2) (2012) 102-108.
- [18] E. Vacin, F. Went, Some pH changes in nutrient solutions. *Bot. Gaz.* 110 (1949) 605–613.
- [19] S. Aktar, K.M. Nasiruddin, K. Hossain, Effects of different media and organic additives interaction on *in vitro* regeneration of *Dendrobium* orchid. *J. Agric. Rural Dev.* 6(1-2) (2008) 69-74.
- [20] S. Kaur, K.K. Bhutani, Organic growth supplement stimulants for *in vitro* multiplication of *Cymbidium pendulum* (Roxb.) Sw. *Hort. Sci. (Prague)* 39(1) (2012) 47–52.

-
- [21] K. Obsuwan, C. Thepsithar, An effect of organic supplements on stimulating growth of *Vanda* and *Mokara* seedlings in tissue culture. *Int. J. Biol. Biomolec. Agric. Food Biotechnol. Eng.* 8(7) (2014) 696-698.
- [22] D. David, R. Jawan, H. Marbaw, J.A. Gansau, Organic additives improves the *in vitro* growth of native orchid *Vanda helvola* Blume. *Not. Sci. Biol.* 7(2) (2015) 192-197.
- [23] Y.S.Huh, J.K. Lee, S.Y. Nam, K.Y. Paek, G.U. Suh, Improvement of asymbiotic seed germination and seedling development of *Cypripedium macranthos* Sw. with organic additives. *J. Plant Biotechnol.* 43 (2016) 138-145.
- [24] J.W.H. Yong, L. Ge, Y.F. Ng, S.N. Tan, The chemical composition and biological properties of coconut (*Cocos nucifera* L.) water. *Molecules* 2009, 14, 5144-5164; doi:10.3390/molecules 14125144.
- [25] P. Gnasekaran, R. Poobathy, M. Mahmood, M.R. Samian, S. Sreeramanan, Effect of complex additives on improving the growth of PLBs of *Vanda* Kasem's Delight. *Aus. J. Crop Sci.* 6(8) (2012) 1245-1248.
- [26] N. Nambiar, C.S. Tee, M. Maziah, Effects of organic additives and different carbohydrate sources on proliferation of protocorm-like bodies in *Dendrobium* Alya Pink. *Plant Omics J.* 5(1) (2012) 10-18.
- [27] M.O. Islam, M. Akter, A.K.M.A. Prodhon, Effect of potato extract on in vitro seed germination and seedling growth of local *Vanda roxburgii* orchid. *J. Bangladesh Agric. Univ.* 9(2) (2011) 211-215.
- [28] Md.A. Baque, Y.K. Shin, T. Elshmari, E.J. Lee, K.P. Paek, Effect of light quality, sucrose and coconut water concentration on the microporpagation of *Calanthe* hybrids ('Bukduseong' × 'Hyesung' and 'Chunkwang' × 'Hyesung'). *Aus. J. Crop Sci.* 5(10) (2011) 1247-1254.
- [29] P. Sinha, S.K. Roy, Regeneration of an indigenous orchid, *Vanda teres* (Roxb.) Lindl. Through in vitro culture. *Plant Tiss. Cult.* 14 (2004) 55-61.
- [30] M. Minea, C. Piluek, T.A. Menakani, S. Tantiwiwat, A study on seed germination and seedling development of *Spathoglottis kimballiana* Bl. orchids. *Kasetsart J. (Nat. Sci.)* 38 (2004) 141-156.