Synthesis, Spectroscopic Characterization and Biological Applications of Novel Zinc Complexes with Schiff Bases

MEHRINE Rehman^{1,a,*}, SAQIB Ali², KHURRAM Shahzad Munawar²

¹Centres of Science and Applied Technologies (CESAT), Islamabad, Pakistan

²Department of Chemistry, Quaid-i-Azam University, Islamabad, 45320, Pakistan

^armehrine@gmail.com

Keywords: Schiff Base, Diabetes Mellitus, Anticancer, Antitumor, DNA Binding Activity, UV Spectroscopy, Antibacterial, Antifungal, Disc Diffusion Method.

Abstract. Schiff base ligands, having nitrogen and oxygen donor sites and their zinc(II) complexes have been prepared by using methanol as a solvent. Purity of the compounds were tested by TLC and characterized by FT-IR, multinuclear NMR (¹H and ¹³C). FT-IR spectra suggested the binding of precursors and their chelation with zinc(II) moieties by showing characteristics peaks in particular regions. ¹H-NMR also approves the synthesis of compounds by showing characteristics peaks of azomethine proton (HC=N) and phenolic proton (-OH). On complexation phenolic proton was disappeared while a down field shift was observed in azomethine proton. ¹³C-NMR data further supported the formation of compounds by displaying and shifting of characteristic peaks of azomethine carbon (HC=N). The DNA binding ability of all the synthesized ligands was studied by UV-Visible spectroscopy. A hypochromic affect was observed showing that ligands interacted with DNA by intercalation. The results implied that these synthesized compounds can be employed for the formulation of Anticancer and Antitumor drugs in future with less toxic side effects to normal cells (unlike toxic drugs presently used). Moreover they were found to have antifungal and antibacterial activity by Disc Diffusion method. Zinc containing complexes had been investigated a lot recently for the treatment of Diabetes Mellitus. Thus, these synthesized compounds are potential drug candidates for research if explored.

Introduction

Many of the diseases had been treated in ancient ages by using variety of metal comprising inorganic compounds. The advancement in medicinal usage of inorganic compounds started in early twenties for the cure of diseases.

Table 1 Established and potent zinc pharmaceuticals

Element	Year	Treatment	Compound
Zn	2002	Stomach ulcer, Diabetes mellitus	Zinc complexes
	2003	UV-induced dermatitis	Zinc complexes
	2007	Metabolic syndrome & Diabetes	Zinc-thioalloxan- <i>N</i> -methyl Zinc-dithiocarbamate
	2008	Wilson disease	Zinc-acetate

Deficiency of zinc in kids is responsible for stunt growth, delay of sexual maturity, weakening of immune system, diarrhea and many further related diseases that lead to increase mortality rate in infants especially in developing countries [1]. There are many biological applications of zinc like it is necessary for development and growth. Presently, antidiabetic activities of zinc are very significant. Zinc and diabetes mellitus (DM) are related at many steps in the process of metabolism within a cell [2].

Fig. 1 Biological uses of zinc

In 1992, it was found that it controls blood glucose level as that of insulin by regulating rat adipocytes lipogenesis followed by *in-vivo* observation of insulin mimetic effects of orally administrated zinc chloride in Streptozotocin-induced type (STZ) diabetic rats and mice [3]. One of the major drawbacks of ZnCl_2 was its low bioavailability, so sufficiently high amount of dose or treatment for the long time (more than 8 weeks) is required. That's why in order to avoid these problems the zinc metal complexes were suggested. In 2002, first orally active antidiabetic zinc complex was discovered. Since then, an extensive range of insulin mimetic zinc complexes having diverse coordination sites around zinc has been suggested [4].

Zinc stimulates lipogenesis and transport of glucose transport in adipocyte tissues; its administration *via* diet decreases high blood sugar level in artificially diabetic mice. Medical investigations also disclosed that there is a connection amongst deficiency of zinc and diabetes mellitus [5]. Few years back, in animals with outstanding antidiabetic and anti-metabolic syndrome properties of the zinc containing complexes have been investigated as the compounds have excellent hypoglycemic abilities on oral administration [6]. Investigations on zinc as a model role in the treatment of diabetes mellitus have been conducted on the diabetic models of animals. The investigations revealed that increased ingestion of zinc containing food remarkably decreased the harshness of type 1 diabetes mellitus.

Normally Schiff bases are generally synthesized by the condensation reactions of carbonyls and amines under different conditions and by using variety of solvents with the elimination of water molecules. The excellent chelation ability of Schiff bases along other properties like easy synthesis and flexibility in changing the chemical surroundings to azomethine specie ($\text{HC}=\text{N}$). Generally, transition metal complexes of Schiff bases are synthesized by reacting them with salts of metals.

DNA is genetic material for all higher organisms.

The basic components of DNA are.

- Deoxyribose sugar having five carbon atoms in the form of cyclic ring
- Phosphate group
- Nitrogenous bases, adenine (A), guanine (G), cytosine (C) and thymine (T)

A double hydrogen bond is developed between A and T, while G and C makes a triple hydrogen bond [7,8]. DNA is the specific point of action of almost all anticancer/antitumor drugs in order to regulate the uncontrolled cell division. These anti-tumor drugs make an adduct with DNA by blocking the particular active sites of DNA which are responsible for replication and elongation of chain. DNA can interact covalently to metal center through oxygen atom of sugar phosphate backbone or to the nitrogen atom of nitrogenous bases, or non-covalently through intercalation or hydrogen bonding. The metal complexes or drugs that interact with DNA can inhibit the processes

related to DNA, such as replication and transcription by forming drug DNA adducts or by distortions of DNA structure leading to apoptosis or cell death.

Experimental

The chemicals used in the present study are of analytical grade and used without additional purification. Zinc salt e.g. $\text{Zn}(\text{CH}_3\text{COO})_2 \cdot 2\text{H}_2\text{O}$ was purchased from Sigma Aldrich and the precursors for the synthesis of ligands were of Acros organics.

Solvents were purified where needed, before use according to the standard procedure [9]. Gallenkamp (Japan) was used to find out the melting point of synthesized compounds with the help of capillary tubes. Infra-red spectral studies, in the range of 4000 to 250 cm^{-1} were carried out on a Nicolet-6700 FT-IR spectrophotometer; Proton and Carbon-13 NMR spectra in deuterated solvents. The electronic absorption spectra were taken on a Shimadzu 1800 UV-Visible Spectrophotometer.

Ligands were synthesized by refluxing stoichiometric amounts of 2-hydroxy-1-naphthaldehyde, an aldehyde with p-toluidine (HL^1) and 2-aminophenol (HL^2) separately in absolute ethanol for 3-4 hours.

(1)

where; X=H, Y=CH₃ for HL^1 and X=OH, Y=H for HL^2

Generalized chemical procedure for the synthesis of ligands HL^1 and HL^2 . Schiff bases, HL^3 and HL^4 were prepared by the condensation of equimolar (1:1) amounts of 5-chloro-2-hydroxybenzaldehyde with 2-aminophenol (HL^3) and ethyl 4-aminobenzoate (HL^4), separately, in absolute ethanol (100 mL). Completion of the reaction was monitored by TLC. The crystalline products were separated by filtration, purified by washing with absolute ethanol and finally dried under vacuum. Zinc(II) complexes 1-4 were synthesized by mixing $\text{Zn}(\text{CH}_3\text{COO})_2 \cdot 2\text{H}_2\text{O}$ with HL^1 and HL^4 in 1:2 ratio and HL^2 and HL^3 with 1:1, using ethanol (100 ml) as a solvent.

(2)

The pure bacterial cultures obtained were maintained in the medium in slants and petri plates. For inoculums preparation, 13 g of nutrient broth was suspended in one liter of distilled water, homogenously mixed and autoclaved for 15 minutes at 121°C. Then 10 μ L of pure culture of a bacterial strain was added to 100 mL of freshly prepared nutrient broth medium. The fungal strains (*Alternaria alternata*, *Ganoderma lucidum*, *Aspergillus niger* and *Penicillium notatum*) were cultured overnight at 28°C using potato dextrose agar. Antimicrobial activities were measured by using disc diffusion method [12]. 2.8 g nutrient agar (for antibacterial activities) or 3.9 g potato dextrose agar (for antifungal activities) was suspended in 100 mL distilled water and sterilized by autoclaving at 121°C for 15 minutes. It was then mixed well with 100 μ L inoculums and was poured in sterilized petri plates. Clear zones of inhibition formed around the discs were measured in millimeters using a zone reader [13]. The zone inhibition was calculated by using the following equation.

$$\% \text{ Growth inhibition} = 100 - \left(\frac{\text{Linear growth in test sample (mm)}}{\text{Linear growth in control (mm)}} \times 100 \right) \quad (6)$$

Results and Discussions

Schiff base ligands and their zinc complexes are synthesized in excellent yield. All the synthesized products are in solid state, having sharp melting points. The synthesized products have been characterized by FT-IR and multinuclear NMR (^1H and ^{13}C) spectroscopy.

The FT-IR bands of complexes were assigned by comparing with the free ligand data and their zinc complexes. All the synthesized ligands have two donor sites which are (i) phenolic oxygen (OH) and (ii) azomethine nitrogen (HC=N). Azomethine group of all the Schiff base ligands gives a strong and sharp peak around 1615-1595 cm^{-1} , which is shifted to lower value around 1613-1580 cm^{-1} in the complexes advocates its coordination to the metal center. The typical phenolic $\nu(\text{OH})$ peak for all ligands was observed at 3443-3455 cm^{-1} is wiped out on complexation presenting the attachment of phenolic oxygen with metal ion. This chelation is further reinforced by the entrance of new peaks at 533-545 cm^{-1} in complexes due to $\nu(\text{Zn-O})$ [14].

In the ^1H -NMR spectra of Schiff base ligands and complexes phenolic protons (-OH) usually appear very down field because of hydrogen bonding and are vanished upon complexation. The signals of azomethine protons are shifted down field giving clear indication/evidence of complex formation. ^{13}C -NMR spectral data of the ligands and complexes were recorded in deuterated solvents. The signals of the azomethine carbon (HC=N) in the ligands are observed at 178.1-160.4 ppm, which after complexation with zinc are shifted down field. Another significant downfield shift upon complexation was observed for the phenolic carbon for complexes of ligand. These peaks shift are good evidence for complexation of ligand with metal atom.

Fig. 2 ^1H -NMR spectrum of HL⁴

The interaction of synthesized ligands HL²-HL⁴ with SS-DNA has been investigated by UV-Visible spectroscopy by monitoring the changes in absorption properties of DNA or drug molecules. The UV-Visible titrations of ligands in absence and in presence of SS-DNA have been studied. Negative values of ΔG showed the spontaneity of the interaction of the drug with DNA.

Fig. 3 Absorption spectra of the compounds HL² in the absence (a) and presence of 10 μM (b), 19 μM (c), 27 μM (d), 35 μM (e), 42 μM (f), 48 μM (g), 54 μM (h) 59 μM (i), 64 μM (j) and 69 μM (k) of DNA

Fig. 4 Absorption spectra of the compounds HL³ in the absence (a) and presence of 10 μM (b), 19 μM (c), 27 μM (d), 35 μM (e), 42 μM (f), 48 μM (g) and 54 μM (h) of DNA

Thus, these synthesized compounds interact with DNA leading to improper functioning and cleavage of the DNA double helix structure. The binding constant values are in the range of 5.38×10^3 to $18.4 \times 10^3 \text{ M}^{-1}$. The negative values of ΔG showed the spontaneity of complex-DNA binding. The moderate binding implies less toxic side effects to normal cells and consequently these synthesized compounds can be employed for the formulation of Anticancer & Antitumour drugs with less toxic side effects in future.

The synthesized ligands HL²-HL⁴ were screened to investigate their *in vitro* behavior against strains of bacteria (*Escherichia coli*, *Bacillus subtilis*, *Staphylococcus aureus* and *Pasturella multocida*) by Disc Diffusion method. Streptomycin was used as the positive controls, while DMSO was negative control. The activities were maximum for strain *Pasturella multocida*, while they were least active against *Escherichia coli*. Ligand HL³ was most active among against all strains of bacteria. More bulky structures results in greater lipophilicity and hence their greater diffusion through the microbial cell membrane thus ultimately causing the greater sensitivity of the

microbes. The difference in the efficiency of various bioactive compounds towards organisms relies upon their permeability through cell membrane [15].

Table 2 Comparison of antibacterial activities of ligands HL²-HL⁴

Compounds	Average zone of inhibition (mm)			
	<i>E. coli</i>	<i>B. subtilis</i>	<i>S. aureus</i>	<i>P. multocida</i>
Streptomycin	30 ^a ±0.26	30 ^a ±0.19	30 ^a ±0.23	30 ^a ±0.29
HL ²	20 ^{bc} ±0.14	19 ^{bc} ±0.13	18 ^c ±0.13	21 ^{bc} ±0.16
HL ³	20 ^{bc} ±0.13	20 ^{bc} ±0.11	19 ^{bc} ±0.15	21 ^{bc} ±0.12
HL ⁴	17 ^c ±0.14	18 ^c ±0.15	19 ^{bc} ±0.15	20 ^{bc} ±0.15

Antibacterial values are mean ± S.D of samples analyzed individually in triplicate at $p < 0.1$; Different letters in superscripts indicate significant differences. a = maximum activity, b = intermediate activity, c = minimum activity, ab = activity between maximum and intermediate and bc = activity between intermediate and minimum.

The maximum antifungal action, from all the studied ligands was observed by HL³. This might be due to more electronegative nature of the chloro group attached to the benzene ring. Also its insulin mimetic properties can be further explored. These complexes are potent drug candidates for further research.

Table 3 Comparison of antifungal activities of ligands HL²-HL⁴

Compounds	Average zone of inhibition (mm)			
	<i>A. alternata</i>	<i>G. lucidum</i>	<i>A. niger</i>	<i>P. notatum</i>
Fluconazole	38 ^a ±0.41	41 ^a ±0.52	40 ^a ±0.36	35 ^a ±0.27
HL ²	20 ^c ±0.12	22 ^{bc} ±0.15	18 ^c ±0.15	20 ^c ±0.10
HL ³	22 ^{bc} ±0.21	25 ^{bc} ±0.22	22 ^{bc} ±0.23	26 ^{bc} ±0.25
HL ⁴	24 ^{bc} ±0.08	21 ^c ±0.34	23 ^{bc} ±0.28	25 ^{bc} ±0.16

letters in superscripts indicate significant differences. a = maximum activity, b = intermediate activity, c = minimum activity, ab = activity between maximum and intermediate and bc = activity between intermediate and minimum.

Conclusion

New azomethine bearing ligands were synthesized, while characterized by FT-IR and NMR spectroscopy. FT-IR confirmed the synthesis of ligands depicting CH=N peaks; which are characteristic of azomethine moiety. NMR spectra gave CH=N signals verified the binding of precursor for ligand synthesis. All the synthesized complexes spontaneously tied with DNA, shown by negative ΔG , impairing DNA structure, via UV-Visible spectroscopy; which made them potential anticancer antitumor drugs to further research work. Their moderate binding constants/affinity meant they were less toxic to normal cells in curing cancers or tumors. Besides others HL² especially had remarkable antibacterial and antifungal activity so these synthesized complexes could be investigated for future drugs.

References

- [1] K.M. Hambidge, N.F. Krebs, Zinc deficiency: a special challenge, J. Nutr., 137 (2007) 1101-05.
- [2] H. Sakurai, Y. Kojima, Y. Yoshikawa, K. Kawabe, H. Yasui, Antidiabetic vanadium (IV) and zinc (II) complexes, Coord. Chem. Rev., 226 (2002) 187-198.
- [3] A. Shisheva, D. Gefel, Y. Shechter, Insulinlike effects of zinc ion in vitro and in vivo: preferential effects on desensitized adipocytes and induction of normoglycemia in streptozocin-induced rats, Diabetes, 41 (1992) 982-988.

-
- [4] H. Sakurai, A. Katoh, Y. Yoshikawa, Chemistry and biochemistry of insulin-mimetic vanadium and zinc complexes. Trial for treatment of diabetes mellitus, *Bull. Chem. Soc. Jpn.*, 79 (2006) 1645-1664.
- [5] M. K. Song, M.J. Rosenthal, S. Hong, D. M. Harris, I. Hwang, I. Yip, M.S. Golub, M.E. Ament, V.L.W. Go, Synergistic antidiabetic activities of zinc, cyclo (his-pro), and arachidonic acid, *Metabolism*, 50 (2001) 53-59.
- [6] Y. Yoshikawa, Y. Adachi, H. Sakurai, A new type of orally active anti-diabetic Zn (II)-dithiocarbamate complex, *Life Sci.*, 80 (2007) 759-766.
- [7] M. Sirajuddin, S. Ali, A. Badshah, Drug–DNA interactions and their study by UV–Vis., fluorescence spectroscopies and cyclic voltametry, *J. Photochem. Photobiol. B: Biol.* 124 (2013) 1-19.
- [8] V. González-Ruiz, A. I. Olives, M. A. Martín, P. Ribelles, M. T. Ramos, J. C. Menéndez, An overview of analytical techniques employed to evidence drug-DNA interactions. Applications to the design of genosensors, *Biomed. Eng. Trend. Res. Technol.*, 32 (2011) 65-90.
- [9] W.L.F. Armarego, C.L.L. Chai, Purification of laboratory chemicals, (2013), 7th Ed; Elsevier (USA).
- [10] S. Dey, S. Sarkar, H. Paul, E. Zangrando, P. Chattopadhyay, Copper (II) complex with tridentate N donor ligand: Synthesis, crystal structure, reactivity and DNA binding study, *Polyhedron* 29 (2010) 1583-1587.
- [11] S. Shujha, A. Shah, N. Muhammad, S. Ali, R. Qureshi, N. Khalid, A. Meetsma, Diorganotin (IV) derivatives of ONO tridentate Schiff base: Synthesis, crystal structure, in vitro antimicrobial, anti-leishmanial and DNA binding studies, *Eur. J. Med. Chem.* 45 (2010) 2902-2911.
- [12] K. Rizwan, M. Zubair, N. Rasool, M. Riaz, M. Zia-Ul-Haq, V. de Feo, Phytochemical and biological studies of *Agave attenuata*, *Int. J. Mol. Sci.* 13 (2012) 6440-6451.
- [13] N. R. Bhalodia, V. Shukla, Antibacterial and antifungal activities from leaf extracts of *Cassia fistula* L.: An ethnomedicinal plant, *J. Adv. Pharm. Technol. Res.* 2 (2011) 104-109.
- [14] T. Yu, W. Su, W. Li, Z. Hong, R. Hua, M. Li, B. Chu, B. Li, Z. Zhang, Z. Z. Hu, Synthesis, crystal structure and electroluminescent properties of a Schiff base zinc complex, *Inorg. Chim. Acta* 359 (2006) 2246-2251.
- [15] C.D. Sheela, C. Anitha, P. Tharmaraj, D. Kodimunthri, Synthesis, spectral characterization, and antimicrobial studies of metal complexes of the Schiff base derived from [4-amino-N-guanylbenezene sulfonamide] and salicylaldehyde, *J. Coord. Chem.*, 63 (2010) 884-893.